

of these species could have developed from pelagic larval forms which drifted into their areas of capture. Their small size and capture in the warmer months (*O. trachypomus* in June, August, and September and *C. parrai* in July) lend credence to this hypothesis.

Increased collecting in progress off the Atlantic coast of the United States will likely reveal that many species now considered sub-tropical or tropical have far more extensive northerly ranges—as strays or random wanderers, as migrants, and as indigenous members of the fauna.

LITERATURE CITED

- BAILEY, REEVE M., et al. 1960. A list of common and scientific names of fishes from the United States and Canada. Am. Fish. Soc., Spec. Publ., no. 2, pp. 1-102.
- BEEBE, WILLIAM, AND JOHN TEE-VAN. 1933. Field book of the shore fishes of Bermuda. G. P. Putnam's Sons, 337 pp., illus.
- BRIGGS, JOHN C. 1958. A list of Florida fishes and their distribution. Bull. Florida State Mus., vol. 2, no. 8, pp. 223-318.
- COLLINS, RICHARD A., AND ROBERT E. SMITH. 1959. Occurrence of butterfly fish in Mississippi Sound. Copeia, no. 3, p. 252.
- FEDDERN, HENRY A. 1963. Color pattern changes during growth of *Bodianus pulchellus* and *B. rufus* (Pisces: Labridae). Bull. Mar. Sci. Gulf and Caribbean, vol. 13, no. 2, pp. 224-241.
- GIBBS, ROBERT H., JR., AND BRUCE B. COLLETTE. 1959. On the identification, distribution, and biology of the dolphins, *Coryphaena hippurus* and *C. equiselis*. Bull. Mar. Sci. Gulf and Caribbean, vol. 9, no. 2, pp. 117-152.
- HOWELL RIVERO, LUIS. 1941. "*Corniger spinosus*" Agassiz: nueva especie para Cuba y algunas consideraciones acerca de la misma. Torreia, no. 6, pp. 3-7.
- HUBBS, CARL L. 1963. *Chaetodon aya* and related deep-living butterflyfishes; their variation, distribution and synonymy. Bull. Mar. Sci. Gulf and Caribbean, vol. 13, no. 1, pp. 133-192.
- JORDAN, DAVID STARR, BARTON WARREN EVERMANN, AND HOWARD WALTON CLARK. 1930. Check list of the fishes and fishlike vertebrates of North and Middle America north of the northern boundary of Venezuela and Colombia. Rept. U. S. Comm. Fish., 1928, App. X., pp. 1-670.

MCKENNEY, THOMAS W. 1959. A contribution to the life history of the squirrel fish *Holocentrus vexillarius* Poey. Bull. Mar. Sci. Gulf and Caribbean, vol. 9, no. 2, pp. 174-221.

U. S. Bureau of Commercial Fisheries Biological Laboratory, Brunswick, Georgia. Contribution number 76.

Quart. Jour. Florida Acad. Sci. 27(4) 1964 (1965)

HYPERTENSIVE EFFECT OF *LATRODECTUS* VENOMS

JOHN D. MCCRONE AND ROGER J. PORTER

THE venoms of several subspecies of the black widow spider, *Latrodectus mactans*, have a hypertensive effect on the mammalian systemic arterial pressure. Troise (1928), Sampayo (1944), Cicardo (1954) and Calvo et al. (1957) demonstrated this with the venom of *L. mactans mactans*, Shapiro et al. (1939) with the venom of *L. mactans indistinctus*, and Bettini and Toschi-Frontali (1960) with the venom of *L. mactans tredecimguttatus*.

This paper presents the results of a study that was undertaken to determine whether the venoms of three other species of *Latrodectus*, *L. variolus*, *L. bishopi*, and *L. geometricus*, have a similar hypertensive effect. This study is part of a more extensive investigation of the comparative toxicology and biochemistry of the venoms of the North American *Latrodectus*.

MATERIALS AND METHODS

Lyophilized venom gland extract reconstituted with physiological saline was used in all experiments. The method of preparation of this extract is given in a previous paper (McCrone, 1964). The glands were taken from mature female *L. mactans mactans*, *L. variolus*, *L. bishopi*, and *L. geometricus* collected in the state of Florida.

The 2.1-3.8 kg rabbits used in this study were anesthetized with intravenous Nembutal (Abbot). The dose for each rabbit was determined individually by observing the eyelid reflex, the average being about 5-6 cc of a 12.5 mg/cc solution. The rabbits were then heparinized with intravenous Panheparin (Abbot). The femoral artery was exposed and cannulated with a small polyethylene tube which was connected to a recording mercury manometer by means of a hypodermic extension tube. Locke's solution containing a small amount of the anticoagulant dye chlorazol fast pink was used in the connecting tube. After normal mean arterial pressure was established and maintained, various dosages of venom were injected into the ear veins of the rabbits and the changes in the mean arterial pressure were recorded on smoked kymograph paper.

TABLE I
Effect of four *Latrodectus* venoms on mean systemic arterial pressure of rabbits

Rabbit	Venom	Dose (mg/kg)	Mean Pressure before Injection (mm Hg)	Latent Period (min.)	Max. Mean Pressure after Injection (mm Hg)	Net Increase in Mean Pressure (mm Hg)	Time (min.) to reach Max. Mean Pressure
1	mactans	0.033	87	0:30	137	50	1:10
2	mactans	0.080	111	1:00	122	11	1:30
3	mactans	0.158	106	0:50	147	41	4:50
4	variolus	0.067	79	1:50	106	27	6:20
5	variolus	0.190	116	1:00	167	51	3:30
6	variolus	0.214	112	1:20	127	15	41:50
7	bishopi	0.067	115	0:50	119	4	2:00
8	bishopi	0.178	67	0:27	89	22	4:10
9	bishopi	0.232	103	0:58	160	57	11:20
10	geometricus	0.035	90	2:20	103	13	5:20
11	geometricus	0.083	89	0:55	119	30	15:25
12	geometricus	0.261	96	1:55	130	34	6:25